

FOREST TYPE: NORTHERN HARDWOODS (BEECH)

SPECIES LIST

Asterisk denotes Species of Greatest Conservation Need. Hyperlinks take you to recommendations specific to the species.

Reptiles and Amphibians

[Five-lined skink](#)*

Northern redbelly snake

[Eastern box turtle](#)*

Northern black racer*

Birds

Common goldeneye

Hooded merganser

Common merganser

[Sharp-shinned hawk](#)*

[Northern goshawk](#)*

[Broad-winged hawk](#)*

[American kestrel](#)*

[American woodcock](#)*

Barred owl

Northern saw-whet owl

[Whip-poor-will](#)*

Downy woodpecker

Hairy woodpecker

Pileated woodpecker

Willow flycatcher

Least flycatcher

[Eastern kingbird](#)*

[White-eyed vireo](#)*

Yellow-throated vireo

Warbling vireo

Blue jay

[Field sparrow](#)*

Black crowned chickadee

White-breasted nuthatch

Eastern bluebird

[Veery](#)*

Swainson's thrush

Hermit thrush

[Wood thrush](#)*

Tennessee warbler

Northern parula

Chestnut-sided warbler

[Black-throated blue warbler](#)*

Black-throated green warbler

[Black-and-white warbler](#)*

[Golden-winged warbler](#)*

American redstart

Ovenbird

[Mourning warbler](#)*

Common yellowthroat

[Yellow-breasted chat](#)*

[Scarlet tanager](#)*

[Rose-breasted grosbeak](#)*

Mammals

Smoky shrew

Snowshoe hare

White-footed mouse

Southern red-backed vole

Woodland jumping mouse

[Red bat](#)*

[Silver-haired bat](#)*

[Little brown bat](#)*

[Hoary bat](#)*

[Indiana bat](#)*

[Northern long-eared bat](#)*

[Tri-colored bat](#)*

Porcupine

Red fox

Gray fox

Fisher

RECOMMENDATIONS

These recommendations are designed to optimize wildlife habitat conditions within this forest type. Other silvicultural options may apply, but they won't necessarily optimize potential habitat conditions for the full range of wildlife species that can occupy this type.

- Use even-aged management. Openings can range from 5 to 50 acres.
- Use a 120-year rotation age with entries every 15 to 20 years.
- Let 10 percent of the area in this type age to 175 years before rotating.
- Favor yellow birch to grow where possible.
- In larger cut blocks (greater than 10 acres), leave patches of uncut trees, about $\frac{1}{3}$ acre for every 10 acres of clear cut.
- Avoid entry during nesting season—April to June.
- Whole-tree harvest or cut-to-length is preferred.